arXiv:1201.2020v2 [math.CO] 10 Apr 2012

THE m-TAMARI LATTICES ARE EL-SHELLABLE

HENRI MÜHLE

Abstract. In this article we prove that the m-Tamari lattices $\mathcal{T}_n^{(m)}$ recently introduced by Bergeron and Préville-Ratelle are EL-shellable in the non-pure sense of Björner and Wachs. Moreover, we characterize the intervals of $\mathcal{T}_n^{(m)}$ according to their topological properties.

1. Introduction

The classical Tamari lattices $\mathcal{T}_n$, see [8], are a well-studied member of the large group of Catalan objects. Bergeron and Préville-Ratelle [1] have recently generalized this class of lattices to m-Tamari lattices $\mathcal{T}_n^{(m)}$ in order to calculate the graded Frobenius characteristic of the space of higher diagonal harmonics. Along with this generalization, they proposed a realization of these lattices via m-Dyck paths, analogously to the realization of the classical Tamari lattices via classical Dyck paths. The main purpose of this article is to prove that the m-Tamari lattices possess a certain order-theoretic property, called EL-shellability, which is recalled in Section 2.2. This is the statement of our main theorem.

Theorem 1.1. *The m-Tamari lattices $\mathcal{T}_n^{(m)}$ are EL-shellable for every positive integer m and n.*

We recall the construction of m-Tamari lattices, as well as the definition of EL-shellability of a partially ordered set (poset) in Section 2. In Section 3 we give an EL-labeling of $\mathcal{T}_n^{(m)}$ by generalizing a construction from [5]. We conclude this article by giving some applications of Theorem 1.1 in Section 4, and we characterize the intervals of $\mathcal{T}_n^{(m)}$ according to their topological properties.

2. Preliminaries

2.1. **Generalized Tamari Lattices.** An *m-Dyck path of height n* is a path from $(0,0)$ to (mn,n) in $\mathbb{N}\times\mathbb{N}$ consisting of steps of the form $(0,1)$ or $(1,0)$, which stays above the line $x=my$. Denote the set of all m-Dyck paths of height n by $\mathcal{D}_n^{(m)}$. It is well-known (see for instance [7]) that the number of m-Dyck paths of height n is counted by the

---

This work was funded by the FWF research grant no. Z130-N13.

Fuss-Catalan number

$$C_n^{(m)} = \frac{1}{mn+1}\binom{(m+1)n}{n}. \tag{1}$$

With any $p \in \mathcal{D}_n^{(m)}$, we can associate a sequence $\alpha_p = (a_1, a_2, \ldots, a_n)$ which satisfies the conditions

$$\begin{aligned} &a_1 \leq a_2 \leq \cdots \leq a_n \quad \text{and} \\ &a_i \leq m(i-1), \quad 1 \leq i \leq n. \end{aligned}$$

The i-th entry of α_p is the smallest non-negative integer at which the corresponding path reaches height i (see Figure 1).



(a) A 4-Dyck path of height 6 with associated sequence $(0, 0, 3, 4, 4, 16)$.



(b) The sequence $(0, 6, 8, 10, 12, 21)$ does not encode a 4-Dyck path.

FIGURE 1. Some Examples of paths and sequences.

For every $i \in \{1, 2, \ldots, n\}$ there is a unique subsequence $(a_i, a_{i+1}, \ldots, a_k)$ of α_p, called *primitive subsequence* that satisfies

$$\begin{aligned} &a_j - a_i < m(j-i), \quad i < j \leq k \quad \text{and} \\ &\text{either} \quad k = n \quad \text{or} \quad a_{k+1} - a_i \geq m(k+1-i). \end{aligned}$$

The dotted line in Figure 1(a) indicates that $(3, 4, 4)$ is the unique primitive subsequence at position 3 in the 4-Dyck path given there.

We construct a covering relation $\lessdot$ on $\mathcal{D}_n^{(m)}$ in the following way: let $p, p' \in \mathcal{D}_n^{(m)}$ such that $\alpha_p = (a_1, a_2, \ldots, a_n)$ satisfies $a_{i-1} < a_i$ for some i. Define

$$\begin{aligned} p \lessdot p' \quad &\text{if and only if} \\ &\alpha_{p'} = (a_1 \ldots, a_{i-1}, a_i - 1, a_{i+1} - 1 \ldots, a_k - 1, a_{k+1}, \ldots, a_n), \end{aligned}$$



FIGURE 2. The lattice of 2-Dyck paths of height 3 and the associated lattice of sequences.

where $(a_i, a_{i+1}, \ldots, a_k)$ is the unique primitive subsequence of α_p at position i. Denote by $\leq$ the reflexive and transitive closure of $\lessdot$. Proposition 4 in [6] implies that the poset $\left(\mathcal{D}_n^{(m)}, \leq\right)$ is a lattice, the *m-Tamari lattice* $\mathcal{T}_n^{(m)}$.

Let $\alpha = (a_1, a_2, \ldots, a_n)$. Whenever we write $\alpha \in \mathcal{T}_n^{(m)}$ in the remainder of this article, we mean $p \in \mathcal{T}_n^{(m)}$ such that α is the sequence associated with p.

Example 2.1. Figure 2 shows the lattice of all 2-Dyck paths of height 3 and the associated lattice of sequences as defined in the previous paragraph.

2.2. **EL-Shellability of Posets.** Initially, this property was introduced for graded posets, in order to create an order-theoretic tool to investigate shellability of posets [2]. Shellability of a poset implies several topological and order theoretical properties, for example Cohen-Macaulayness of the associated order complex. More implications of shellability can for instance be found in [2–5]. In [4], Björner and Wachs generalized EL-shellability to non-graded posets. This is the property of interest in the present article.

Let $(P, \leq)$ be a poset. We call a poset *bounded* if it has a unique minimal and a unique maximal element, denoted by $\hat{0}$ and $\hat{1}$, respectively. Denote by $\mathcal{E}(P)$ the set of all covering relations $p \lessdot q$ in P. Hence, $\mathcal{E}(P)$ corresponds to the set of edges in the Hasse diagram of P. Consider a non-singleton interval $[x, y]$ in P and a chain $c : x = p_0 < p_1 < \cdots < p_s = y$. A chain is called *maximal in* $[x, y]$ if there is no $q \in P$ and no $i \in \{1,2,\ldots,s\}$ such that $p_0 < p_1 < \cdots < p_{i-1} < q < p_i < \cdots < p_s$. For some poset Λ, call a map $\lambda : \mathcal{E}(P) \to \Lambda$ *edge-labeling of* P and let $\lambda(c)$ denote the sequence $\bigl(\lambda(p_0,p_1),\lambda(p_1,p_2),\ldots,\lambda(p_{s-1},p_s)\bigr)$ of edge-labels of c with respect to λ. The chain c is called *rising* if $\lambda(c)$ is a strictly increasing sequence. Moreover, c is called *lexicographically smaller* than another maximal chain $\tilde{c}$ in the same interval if $\lambda(c) < \lambda(\tilde{c})$ with respect to the lexicographic order on Λ^*, the set of words over the alphabet Λ. More precisely, the lexicographic order on Λ^* is defined as $(p_1,p_2,\ldots,p_s) \leq (q_1,q_2,\ldots,q_t)$ if and only if either

$$\begin{aligned} p_i = q_i,&\quad \text{for } 1 \leq i \leq s \text{ and } s \leq t \quad \text{or}\\ p_i < q_i,&\quad \text{for the least } i \text{ such that } p_i \neq q_i. \end{aligned}$$

An edge-labeling of P is called *EL-labeling* if for every interval $[x,y]$ in P there exists a unique rising maximal chain in $[x,y]$, which is lexicographically first among all maximal chains in $[x,y]$. A bounded poset that admits an EL-labeling is called *EL-shellable*.

3. EL-Shellability of $\mathcal{T}_n^{(m)}$

Björner and Wachs have shown in [5, Section 9] that the classical Tamari lattices are EL-shellable. Hence, it is a natural question to ask whether the m-Tamari lattices have the same property. Although Björner and Wachs use slightly different sequences to represent the elements of $\mathcal{T}_n$, their proposed labeling function gives a good intuition on how to generalize their result.

Consider the edge-labeling

$$\begin{aligned} \lambda : \mathcal{E}\bigl(\mathcal{T}_n^{(m)}\bigr) &\to \mathbb{N}\times\mathbb{N}\\ (\alpha,\beta) &\mapsto (j,a_j), \end{aligned} \tag{2}$$

where $\alpha = (a_1,a_2,\ldots,a_n), \beta = (b_1,b_2,\ldots,b_n)$, as well as $j = \min\bigl\{i \in \{1,2,\ldots,n\} \mid a_i \neq b_i\bigr\}$.

Example 3.1. Figure 3 shows the Hasse diagram of $\mathcal{T}_3^{(2)}$ together with the edge-labeling defined in (2).

The main result of this section is stated in the following theorem.

Theorem 3.2. *Let $\mathcal{T}_n^{(m)}$ be the m-Tamari lattice of order n. The edge-labeling given in (2) is an EL-labeling of $\mathcal{T}_n^{(m)}$ with respect to the following order on $\mathbb{N}\times\mathbb{N}$*

$$(i,a_i) \leq (j,a_j) \quad \textit{if and only if} \quad i < j, \quad \textit{or} \quad i = j \textit{ and } a_i \geq a_j. \tag{3}$$

Moreover, there is at most one falling chain in each interval of $\mathcal{T}_n^{(m)}$.

Proof. We need to show that for every interval $[\alpha,\beta]$ in $\mathcal{T}_n^{(m)}$ there exists exactly one rising chain that is lexicographically first among all maximal chains in $[\alpha,\beta]$. Let $\alpha = (a_1,a_2,\ldots,a_n)$ and $\beta = (b_1,b_2,\ldots,b_n)$



FIGURE 3. The diagram of $\mathcal{T}_3^{(2)}$ labeled with the edge labeling as defined in (2).

and consider the set $D = \{j \mid a_j \neq b_j\} = \{j_1, j_2, \ldots, j_s\}$. Let the elements of D be listed in increasing order, namely $j_1 < j_2 < \cdots < j_s$. Let $\rho^{(0)} = \alpha$ and construct $\rho^{(i+1)}$ from $\rho^{(i)}$ by decreasing the values in the primitive subsequence at position j_k of $\rho^{(i)}$ by one, where j_k is the smallest element of D such that the j_k-th entry of $\rho^{(i)}$ is larger than b_{j_k}. By the minimality of j_k it is ensured that $\rho^{(i+1)} \leq \beta$. Since $\mathcal{T}_n^{(m)}$ is a finite lattice, we obtain $\rho^{(t)} = \beta$ after a finite number, say t, of steps. By construction, it is clear that the chain $\alpha = \rho^{(0)} < \rho^{(1)} < \cdots < \rho^{(t)} = \beta$ is rising in the interval $[\alpha, \beta]$.

Let $\rho^{(i)} = (r_1^{(i)}, r_2^{(i)}, \ldots, r_n^{(i)})$. It is guaranteed by construction that $\lambda\big(\rho^{(i)}, \rho^{(i+1)}\big) = \big(j, r_j^{(i)}\big) = \min\Big\{\lambda\big(\rho^{(i)}, \bar{\rho}\big) \mid \big(\rho^{(i)}, \bar{\rho}\big) \in \mathcal{E}\big(\mathcal{T}_n^{(m)}\big)\Big\}$. Since the primitive subsequence of $\rho^{(i)}$ at position j is unique, any other covering relation yields a label $\big(j', r_{j'}^{(i)}\big)$, with $j < j'$. By following such a chain upwards, we will eventually encounter an edge (σ, τ) such that the corresponding edge label is (j, s_j), where $\sigma = (s_1, s_2, \ldots, s_n)$. This creates a descent in such a chain. Hence, the constructed chain is the unique rising chain in $[\alpha, \beta]$ and is lexicographically first.

Now consider the set $D' = \{j \mid a_j \neq b_j \text{ and } a_j \geq a_{j-1} + m\} = \{j_1, j_2, \ldots, j_t\}$. This means that $k \in D$ implies $a_k \neq b_k$ and there is no primitive subsequence of α at position $i < k$, which contains a_k. Similarly to the previous paragraph, we can see that if there exists a falling chain $\alpha^{(0)} < \alpha^{(1)} < \cdots < \alpha^{(t)}$ in $[\alpha, \beta]$, it must have the sequence of edge-labels

$$\big(j_t, a_{j_t}^{(0)}\big), \big(j_{t-1}, a_{j_{t-1}}^{(1)}\big), \ldots, \big(j_1, a_{j_1}^{(t-1)}\big),$$

since each of the values $a_{j_1}, a_{j_2}, \ldots, a_{j_t}$ must be decreased along any maximal chain in $[\alpha, \beta]$ at least once. Hence, the given chain is the only possible falling chain. ☐

Proof of Theorem 1.1. This follows by definition from Theorem 3.2. ☐

Remark 3.3. Proposition 4 in [6] states that $\mathcal{T}_n^{(m)}$ is an interval in $\mathcal{T}_{mn}^{(1)}$. Bearing this in mind, the statement of Theorem 1.1 can already be deduced from [5, Theorem 9.2]. (By definition, an interval of an EL-shellable poset is EL-shellable with respect to the restricted labeling.) However, this has not been done before explicitly. Moreover, our construction does not use the mentioned results and gives a uniform description of the EL-shellability of $\mathcal{T}_n^{(m)}$ in terms of m-Dyck paths.

4. APPLICATIONS

According to [4], Theorem 3.2 has some consequences for the Möbius function of $\mathcal{T}_n^{(m)}$ and the structure of the topological space associated to the intervals of $\mathcal{T}_n^{(m)}$. For an introduction on how to associate a topological space to a poset, we refer to [9].

Corollary 4.1. *Let* $[x, y]$ *be an interval of* $\mathcal{T}_n^{(m)}$ *and* μ *the Möbius function of* $\mathcal{T}_n^{(m)}$*. Then,* $\mu(x, y) \in \{-1, 0, 1\}$*.*

Proof. This result follows immediately from Theorem 3.2 and [4, Theorem 5.7]. ☐

Corollary 4.2. *Every open interval in* $\mathcal{T}_n^{(m)}$ *has the homotopy type of either a point or a sphere.*

Proof. The result follows immediately from Theorem 1.1 and [4, Theorem 5.9]. ☐

We can specify the previous results even more and characterize the spherical intervals. For that purpose, consider $\alpha = (a_1, a_2, \ldots, a_n) \in \mathcal{T}_n^{(m)}$ as well as $D \subseteq \{2, 3, \ldots, n\}$ such that $a_j > a_{j-1}$ for all $j \in D$. Define the numbers

$$\text{(4)} \quad \mathrm{ps}_\alpha(j) = \big|\{i \in D \mid i < j \text{ and } a_j - 1 - a_i < m(j-i) \text{ and } a_k - a_i < m(k-i) \text{ for all } i < k < j\}\big|$$

for all $j \in D$. Consider the sequence $\alpha_j^{\uparrow} \in \mathcal{T}_n^{(m)}$ that arises from α by decreasing the primitive subsequence of α at position $j \in D$ by one. Clearly, $(\alpha, \alpha_j^{\uparrow}) \in \mathcal{E}\big(\mathcal{T}_n^{(m)}\big)$ and $\mathrm{ps}_\alpha(j)$ counts the primitive subsequences of $\alpha_j^{\uparrow}$ at some $i \in D$ with $i < j$ that contain the j-th entry of $\alpha_j^{\uparrow}$.

Theorem 4.3. *Let* $\alpha \leq \beta$ *in* $\mathcal{T}_n^{(m)}$, *where* $\alpha = (a_1, a_2, \ldots, a_n)$ *and* $\beta = (b_1, b_2, \ldots, b_n)$. *Let* $D = \{j \mid a_j \neq b_j \text{ and } a_j > a_{j-1}\}$. *The open interval* (α, β) *has the homotopy type of a* $(|D| - 2)$*-sphere if*

$$a_j - 1 - a_{j-1} < m \quad \textit{implies} \quad b_j - b_{j-1} < m, \quad \textit{and} \tag{5}$$

$$b_j = a_j - 1 - ps_\alpha(j), \tag{6}$$

for all $j \in D$. *Otherwise,* (α, β) *has the homotopy type of a point.*

Proof. Let $\alpha = (a_1, a_2, \ldots, a_n)$ and $\beta = (b_1, b_2, \ldots, b_n)$ such that $\alpha \leq \beta$. We need to show that a falling chain exists in $[\alpha, \beta]$ if and only if the conditions (5) and (6) are satisfied.

Write the set D in the form $D = \{j_1, j_2, \ldots, j_s\}$, where $j_1 < j_2 < \cdots < j_s$. Let $\rho^{(0)} = \alpha$ and construct $\rho^{(i+1)}$ from $\rho^{(i)}$ by decreasing the primitive subsequence at position j_{s-i} of $\rho^{(i)}$ by one. It is clear that the chain $\rho^{(0)} < \rho^{(1)} < \cdots < \rho^{(s)}$ is falling. We also notice that $\mathrm{ps}_{\rho^{(i)}}(j_k) = \mathrm{ps}_\alpha(j_k)$ if $k < s - i$.

First we show that (5) is equivalent to $\rho^{(k)} \leq \beta$ for all $k \in \{1, 2, \ldots, s\}$. Assume that there exists a $k \in \{1, 2, \ldots, s\}$ such that $a_{j_k} - 1 - a_{j_k - 1} < m$ and $b_{j_k} - b_{j_k - 1} \geq m$ and j_k is maximal in D with respect to this property. Consider the element $\tilde{\alpha}^{(0)} = \left(\tilde{a}_1^{(0)}, \tilde{a}_2^{(0)}, \ldots, \tilde{a}_n^{(0)}\right) \in \mathcal{T}_n^{(m)}$ that arises from $\rho^{(s-k)}$ by decreasing the primitive subsequence of $\rho^{(s-k)}$ at position j_k by one. Thus, $\tilde{\alpha}^{(0)} = \rho^{(s-k+1)}$. Construct elements $\tilde{\alpha}^{(i+1)}$ from $\tilde{\alpha}^{(i)}$ by decreasing the value of the primitive subsequence of $\tilde{\alpha}^{(i)}$ at position $j_k - 1$ by one. By assumption, we know that $\tilde{a}_{j_k}^{(i)}$ is contained in the primitive subsequence of $\tilde{\alpha}^{(i)}$ at position $j_k - 1$. Hence, in each such step, we decrease the value of $\tilde{a}_{j_k - 1}^{(i)}$ and $\tilde{a}_{j_k}^{(i)}$ (and possibly some subsequent entries). After a finite number, say t, of steps, we obtain an element $\tilde{\alpha}^{(t)} = (\tilde{a}_1^{(t)}, \tilde{a}_2^{(t)}, \ldots, \tilde{a}_n^{(t)})$, such that $\tilde{a}_{j_k - 1}^{(t)} = b_{j_k - 1}$. Since b_{j_k} is not contained in the primitive subsequence of β at position $j_k - 1$, we have $\tilde{a}_{j_k}^{(t)} < b_{j_k}$, and thus $\beta \leq \tilde{\alpha}^{(t)}$. Certainly, there is an $u \in \{0, 1, \ldots, t-1\}$ such that $\tilde{a}_{j_k}^{(u)} = b_{j_k}$, and hence $\tilde{a}_{j_k - 1}^{(u)} > b_{j_k - 1}$. This implies that $\tilde{a}_{j_k - 1}^{(i)} > b_{j_k - 1}$ for all $i \in \{0, 1, \ldots, u\}$, and we can conclude that $\tilde{\alpha}^{(i)} \not\leq \beta$. Thus, $\rho^{(s-k+1)} \not\leq \beta$. The reverse implication is trivial.

Now we show that (6) is equivalent to the fact that $(\rho^{(k-1)}, \rho^{(k)}) \in \mathcal{E}\left(\mathcal{T}_n^{(m)}\right)$ for all $k \in \{1, 2, \ldots, s\}$. The number $\mathrm{ps}_\alpha(j_k)$ corresponds to the number of primitive subsequences of $\rho^{(1)} = \left(r_1^{(1)}, r_2^{(1)}, \ldots, r_n^{(1)}\right)$ at position $i \in D$, where $i < j_k$ that contain $r_{j_k}^{(1)}$. Hence, along the chain

$\rho^{(1)} < \rho^{(2)} < \cdots < \rho^{(s)}$, the value of $r^{(1)}_{j_k}$ is decreased exactly $\mathrm{ps}_\alpha(j_k)$-times and hence $b_{j_k} = r^{(1)}_{j_k} - \mathrm{ps}_\alpha(j_k)$. By constructing $\rho^{(1)}$, we obtain $r^{(1)}_{j_k} = a_{j_k} - 1$ which implies the claim.

By combining both properties, we obtain that $\rho^{(0)} < \rho^{(1)} < \cdots < \rho^{(s)}$ is indeed a falling chain from α to β. ☐

Corollary 4.4. *Let $\alpha \leq \beta$ in $\mathcal{T}_n^{(m)}$, where $\alpha = (a_1, a_2, \ldots, a_n)$ and $\beta = (b_1, b_2, \ldots, b_n)$. Let $D = \{j \mid a_j \neq b_j$ and $a_j > a_{j-1}\}$ and let μ denote the Möbius function of $\mathcal{T}_n^{(m)}$. Then,*

$$\mu(\alpha,\beta) = \begin{cases} (-1)^{|D|}, & \text{if conditions (5) and (6) hold,}\\ 0, & \text{otherwise.}\end{cases}$$

Corollary 4.5. *Let $\alpha \in \mathcal{T}_n^{(m)}$, where $\alpha = (a_1, a_2, \ldots, a_n)$. Let $D = \{j \mid a_j \neq (j-1)m\}$. Let $D_j = \{i \in D \mid i < j\}$, and let μ denote the Möbius function of $\mathcal{T}_n^{(m)}$. Then,*

$$\mu(\hat{0},\alpha) = \begin{cases} (-1)^{|D|}, & \text{if } a_j = (j-1)m - 1 - |D_j| \text{ for all } j \in D,\\ 0, & \text{otherwise.}\end{cases}$$

Proof. By construction, $\hat{0} = (0, m, 2m, \ldots, (n-1)m)$. Hence, the premise in condition (5) corresponds to $(j-1)m - 1 - (j-2)m = m - 1 < m$ and is always satisfied. Moreover, we have

$$\mathrm{ps}_{\hat{0}}(j) = |\{i \in D \mid i < j\}| = |D_j|,$$

for all $j \in D$. If α satisfies (6), then $a_j = (j-1)m - 1 - \mathrm{ps}_{\hat{0}}(j)$. There are two possibilities: either $j-1 \in D$, or $j-1 \notin D$.

Consider the case that $j-1 \in D$. Then,

$$\begin{aligned} a_j - a_{j-1} &= (j-1)m - 1 - \mathrm{ps}_{\hat{0}}(j) - (j-2)m + 1 + \mathrm{ps}_{\hat{0}}(j-1)\\ &= m - \mathrm{ps}_{\hat{0}}(j) + \mathrm{ps}_{\hat{0}}(j-1)\\ &= m - |D_j| + |D_{j-1}|,\end{aligned}$$

which yields $|D_j| - |D_{j-1}| > 0$ for the conclusion of (5). Since $j-1 \in D$, we know that $D_{j-1} \subsetneq D_j$ and the conclusion of (5) is automatically true if $a_j = (j-1)m - 1 - |D_j|$ for all $j \in D$.

Now let $j-1 \notin D$. Then, $a_{j-1} = (j-2)m$, and we have

$$\begin{aligned} a_j - a_{j-1} &= (j-1)m - 1 - \mathrm{ps}_{\hat{0}}(j) - (j-2)m\\ &= m - 1 - \mathrm{ps}_{\hat{0}}(j)\\ &= m - 1 - |D_j|,\end{aligned}$$

which yields $|D_j| + 1 > 0$ for the conclusion of (5). Since $|D_j| \geq 0$, the conclusion of (5) is automatically true if $a_j = (j-1)m - 1 - |D_j|$ for all $j \in D$, which completes the proof. ☐

Corollary 4.6. *Let* $\alpha \in \mathcal{T}_n^{(m)}$*, where* $\alpha = (a_1, a_2, \ldots, a_n)$*. Let* $D = \{j \mid a_j > a_{j-1}\}$ *and let* μ *denote the Möbius function of* $\mathcal{T}_n^{(m)}$*. Then*

$$\mu(\alpha, \hat{1}) = \begin{cases} (-1)^{|D|}, & \text{if } a_j = ps_\alpha(j) + 1 \text{ for all } j \in D \\ 0, & \text{otherwise.} \end{cases}$$

Proof. It follows from the definition of $\mathcal{T}_n^{(m)}$ that $\hat{1} = (0, 0, \ldots, 0)$. Hence, for an interval $[\alpha, \hat{1}]$ in $\mathcal{T}_n^{(m)}$ condition (5) is trivially true for all $j \in D$ and condition (6) reduces to $a_j = \mathrm{ps}_\alpha(j) + 1$ for all $j \in D$. ☐

In the remainder of this section, we prove that the number of spherical intervals of $\mathcal{T}_n^{(m)}$ involving $\hat{0}$ or $\hat{1}$ is 2^{n-1} respectively.

Proposition 4.7. *Let* $m, n \in \mathbb{N}$*. Let* $\mathcal{S}_n^{(m)}(\hat{0}) = \{p \in \mathcal{T}_n^{(m)} \mid \mu(\hat{0}, p) \neq 0\}$ *and* $\mathcal{S}_n^{(m)}(\hat{1}) = \{p \in \mathcal{T}_n^{(m)} \mid \mu(p, \hat{1}) \neq 0\}$*. Then,*

$$|\mathcal{S}_n^{(m)}(\hat{0})| = 2^{n-1} = |\mathcal{S}_n^{(m)}(\hat{1})|. \tag{7}$$

Consider $p, q \in \mathcal{T}_n^{(m)}$, with associated sequences $\alpha_p = (a_1, a_2, \ldots, a_n)$ and $\alpha_q = (b_1, b_2, \ldots, b_n)$. Define

$$D(p, q) = \big\{i \in \{2, \ldots, n\} \mid a_i \neq b_i\big\}.$$

Now, fix $p \in \mathcal{T}_n^{(m)}$ and $D \subseteq \{2, 3, \ldots, n\}$, and define

$$\mathrm{diff}_D(p) = \{q \in \mathcal{T}_n^{(m)} \mid D(p, q) = D\}. \tag{8}$$

It is immediately clear that for every $p \in \mathcal{T}_n^{(m)}$

$$\mathcal{T}_n^{(m)} = \bigcup_{D \subseteq \{2,3,\ldots,n\}} \mathrm{diff}_D(p),$$

and $\mathrm{diff}_{D_1}(p) \cap \mathrm{diff}_{D_2}(p) = \emptyset$ if $D_1 \neq D_2$.

Lemma 4.8. *Let* $D \subseteq \{2, 3, \ldots, n\}$*. Then,* $\mathit{diff}_D(\hat{0}) \neq \emptyset$*.*

Proof. Let $D \subseteq \{2, 3, \ldots, n\}$. The sequence associated to $\hat{0}$ is $\alpha_{\hat{0}} = (0, m, 2m, \ldots, (n-1)m)$. Consider the indicator function

$$\chi_D : \{2, 3, \ldots, n\} \to \{0, 1\}, \quad i \mapsto \begin{cases} 1, & \text{if } i \in D, \\ 0, & \text{otherwise.} \end{cases}$$

Since $m \geq 1$, it is clear that the sequence $\chi_D(\hat{0}) = \big(0, m - \chi_D(2), 2m - \chi_D(3), \ldots, (n-1)m - \chi_D(n)\big)$ corresponds to an m-Dyck path again. ☐

Denote by $[i, j]$ the interval $\{i, i+1, \ldots, j\}$ for $1 \leq i \leq j \leq n$.

Lemma 4.9. *Let* $i \in \{2, 3, \ldots, n\}$*. Then,* $\mathit{diff}_D(\hat{1}) \neq \emptyset$ *if and only if* $D = [i, n]$ *or* $D = \emptyset$*.*

Proof. By construction, $\alpha_{\hat{1}} = (0,0,\ldots,0)$ is the sequence associated to $\hat{1}$. If $D \subseteq \{2,3,\ldots,n\}$ and $D \neq [i,n]$ for some $2 \leq i \leq n$, then any element in $\mathrm{diff}_D(\hat{1})$ has to correspond to a sequence of the form $(0,b_2,b_3,\ldots,b_{n-1},0)$. By definition it is clear that this encodes an m-Dyck path only in the case $b_2 = b_3 = \cdots = b_{n-1} = 0$. Hence, $D = \emptyset$.

Now let $D = [i,n]$ for some $2 \leq i \leq n$. Consider for instance the sequence

$$\alpha = (\underbrace{0,0,\ldots,0}_{i-1},\underbrace{1,1,\ldots,1}_{n-i+1}).$$

This sequence clearly encodes an m-Dyck path, and hence $\mathrm{diff}_D(\hat{1}) \neq \emptyset$. ☐

Proof of Proposition 4.7. We first consider the case $\mathcal{S}_n^{(m)}(\hat{0})$. Let $D = \{j_1,j_2,\ldots,j_t\} \subseteq \{2,3\ldots,n\}$. Let $\alpha^{(0)} = \hat{0}$ and construct $\alpha^{(i+1)}$ from $\alpha^{(i)}$ by reducing the primitive subsequence of $\alpha^{(i)}$ at position j_{t-i} by one. Let $\alpha^{(i)} = (a_1^{(i)},a_2^{(i)},\ldots,a_n^{(i)})$. It is clear by construction that $a_k^{(i)} = a_k^{(0)}$ for all $k < j_{t-i+1}$. Hence, we obtain a falling chain $\alpha^{(0)} \lessdot \alpha^{(1)} \lessdot \cdots \lessdot \alpha^{(t)}$, where $\alpha^{(t)} \in \mathrm{diff}_D(\hat{0})$. (In fact, $\alpha^{(t)}$ corresponds to $\chi_D(\hat{0})$ as defined in the proof of Lemma 4.8.) It follows from the proof of Theorem 3.2 that $\alpha^{(t)}$ is the only m-Dyck path in $\mathrm{diff}_D(\hat{0})$ with this property. Hence, for every $D \subseteq \{2,3,\ldots,n\}$ we obtain exactly one $p \in \mathrm{diff}_D(\hat{0})$, which yields $|\mathcal{S}_n^{(m)}(\hat{0})| = 2^{n-1}$.

Now we show $|\mathcal{S}_n^{(m)}(\hat{1})| = 2^{n-1}$. Let $D = [i,n]$ for some $2 \leq i \leq n$, let $\bar{D} \subseteq \{i+1,i+2,\ldots,n\}$ with $\bar{D} = \{j_1,j_2,\ldots,j_s\}$, and define $j_0 = i$. Consider $p \in \mathrm{diff}_D(\hat{1})$ with the associated sequence

$$\alpha_p = (\underbrace{0,0,\ldots,0}_{j_0-1},\underbrace{1,1\ldots,1}_{j_1-j_0},\underbrace{2,2,\ldots,2}_{j_2-j_1},\ldots,\underbrace{s+1,s+1,\ldots,s+1}_{n-j_s+1}).$$

Let $\alpha^{(0)} = p$ and construct $\alpha^{(i+1)}$ from $\alpha^{(i)}$ by reducing the value of the primitive subsequence of $\alpha^{(i)}$ at position j_{s-i} by one. We obtain a chain $\alpha^{(0)} \lessdot \alpha^{(1)} \lessdot \cdots \alpha^{(s+1)}$, of length $s+1$ with label sequence $\bigl((j_s,s+1),(j_{s-1},s),\ldots,(j_0,1)\bigr)$. Moreover, $\alpha^{(s+1)} = \hat{1}$. This implies that $\mu(p,\hat{1}) = (-1)^{s+1}$. Hence, the total number of elements $p \in \mathrm{diff}_D(\hat{1})$ is 2^{n-i}, satisfying $\mu(p,\hat{1}) \neq 0$. This implies

$$\begin{aligned} |\mathcal{S}_n^{(m)}(\hat{1})| &= 1 + \sum_{i=2}^{n} 2^{n-i} \\ &= 1 + \sum_{i=0}^{n-2} 2^{i} \\ &= 1 + 2^{n-1} - 1 \\ &= 2^{n-1}. \end{aligned}$$

☐

References

[1] François Bergeron and Louis-François Préville-Ratelle, *Higher Trivariate Diagonal Harmonics via Generalized Tamari Posets* (2011), available at `arXiv:1105.3738`.

[2] Anders Björner, *Shellable and Cohen-Macaulay Partially Ordered Sets*, Transactions of the American Mathematical Society **260** (1980), 159–183.

[3] Anders Björner and Michelle L. Wachs, *On Lexicographically Shellable Posets*, Transactions of the American Mathematical Society **277** (1983), 323–341.

[4] ———, *Shellable and Nonpure Complexes and Posets I*, Transactions of the American Mathematical Society **348** (1996), 1299–1327.

[5] ———, *Shellable and Nonpure Complexes and Posets II*, Transactions of the American Mathematical Society **349** (1997), 3945–3975.

[6] Mireille Bousquet-Mélou, Éric Fusy, and Louis-François Préville-Ratelle, *The Number of Intervals in the m-Tamari Lattices* (2011), available at `arXiv:1106.1498v1`.

[7] Aryeh Dvoretzky and Theodore Motzkin, *A Problem of Arrangements*, Duke Mathematical Journal **14** (1947), 305–313.

[8] Dov Tamari, *The Algebra of Bracketings and their Enumeration*, Nieuw Archief voor Wiskunde **10** (1962), 131–146.

[9] Michelle L. Wachs, *Poset Topology: Tools and Applications*, Geometric Combinatorics, 2007, pp. 497–615.

FAK. FÜR MATHEMATIK, UNIVERSITÄT WIEN, GARNISONGASSE 3, 1090 WIEN, AUSTRIA

E-mail address: `henri.muehle@univie.ac.at`